

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

PCT/ SE 99 / 0 1 6 7 1

09/743852

REC'D 16 DEC 1999

WIPO

PCT

Intyg
Certificate

EJW

SE 99/167-1

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

(71) Sökande A+Science Invest AB, Göteborg SE
Applicant (s)

(21) Patentansökningsnummer 9803276-6
Patent application number

(86) Ingivningsdatum 1998-09-25
Date of filing

Stockholm, 1999-12-07

För Patent- och registreringsverket
For the Patent- and Registration Office

Emma Johnsson

Avgift
Fee

PATENT- OCH
REGISTRERINGSVERKET
SWEDEN

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 666 02 86
08-666 02 86

Ink. t. Patent- och reg.verket

1998 -09- 25

Huvudfaxen Kassar

1

USE OF CERTAIN DRUGS FOR TREATING NERVE ROOT INJURY**DESCRIPTION****5 Technical field**

The present invention relates to the use of a metalloproteinase inhibitor in the preparation of pharmaceutical compositions for the treatment of nerve root injury, as well as a method for treating nerve root injury.

- 10 The object of the present invention is to obtain a possibility to treat nerve root injury, which may turn up as sciatica, by blocking disk related cytokines.

Background of the invention

- 15 Disk herniation is a troublesome disorder, which causes great pain and lost of motility, and thereby lost of productive life. Disk herniation will cause trouble to a varying degree, and the pain may last for a month or in severe cases up to 6 months. Beside the pain sciatica may cause severe handicapping problems, as the one suffering therefrom will feel crippled thereof.

- 20 US-A-5,703,092 discloses the use of hydroxamic acid compounds and carbocyclic acids as metalloproteinase and TNF inhibitors, and in particular in treatment of arthritis and other related inflammatory diseases. No use of these compounds for the treatment of nerve root injuries is disclosed or hinted at.

- 25 US-A-4,925,833 discloses the use of tetracyclines to enhance bone protein synthesis, and treatment of osteoporosis.

- 30 US-A-4,666,897 discloses inhibition of mammalian collagenolytic enzymes by tetracyclines. The collagenolytic activity is manifested by excessive bone resorption, periodontal disease, rheumatoid arthritis, ulceration of cornea, or resorption of skin or other connective tissue collagen.

Neither of these latter two documents mentions nerve root injury or the treatment thereof.

5 It has been noted that cytokines from nucleus pulposus-cells such as TNF- α , interleukin 1- β and interferon- γ and others cause structural and functional damages on closely situated nerves in connection with e.g., a disk herniation. Furthermore the nerve becomes sensitised by these substances to produce pain when they are mechanically deformed.

Description of the present invention

10 It has now surprisingly been shown possible to be able to treat nerve root injuries, or at least alleviate the symptoms of nerve root injuries by using a pharmaceutical composition comprising an therapeutically active amount of a metalloproteinase inhibitor.

15 The therapeutically effective amount is a dosage normally used when using such compounds for other therapeutic uses. Many of these drugs are commercially known registered drugs.

20 Compounds that possess this activity are tetracyclines, such as tetracycline, doxycycline, lymecycline, oxytetracycline, minocycline, dedimethylaminotetracycline, hydroxamic acid compounds, carbocyclic acids and derivatives, cyclosporine, methylprenisolone, thalidomide, lazaroides, pentoxiphylline. These can be present as bases or in the form of addition salts, whichever possesses the best pharmaceutical effect, and best property to be brought into a pharmaceutical suitable composition.

25 The effect of doxycycline has been studied and the methods used and results obtained are disclosed below.

Example

Study design.

30 The effects of nucleus pulposus and various treatments to block TNF- α activity were evaluated in an experimental set-up using immunohistochemistry and nerve conduction velocity recordings.

Summary of background data:

A meta-analysis of observed effects induced by nucleus pulposus reveals that these effects might relate to one specific cytokine, Tumor Necrosis Factor alpha (TNF(α)).

5 Objectives.

To assess the presence of TNF(α) in pig nucleus pulposus cells and to see if block of TNF(α) also blocks the nucleus pulposus-induced reduction of nerve root conduction velocity.

10 Methods

Series-1: Cultured nucleus pulposus-cells were immunohistologically stained with a monoclonal antibody for TNF(α).

Series-2: Nucleus pulposus was harvested from lumbar discs and applied to the sacro-coccygeal cauda equina in 13 pigs autologously. Four pigs received 100 mg of doxycycline intravenously, 5 pigs had a blocking monoclonal antibody to TNF- α applied locally in the nucleus pulposus, and 4 pigs remained non-treated and formed control. Three days after the application the nerve root conduction velocity was determined over the application zone by local electrical stimulation.

20 Results.

Series-1: TNF- α was found to be present in the nucleus pulposus-cells.

Series-2: The selective antibody to TNF- α limited the reduction of nerve conduction velocity, although not statistically significantly to the control series. However, treatment with doxycycline significantly blocked the nucleus pulposus-induced reduction of conduction velocity.

Conclusion.

For the first time a specific substance, Tumor Necrosis Factor-alpha, has been linked to the nucleus pulposus-induced effects of nerve roots after local application. Although the effects of this substance may be synergistic with other similar substances, the data of the present study may be of significant importance for the continued understanding of nucleus pulposus'

biologic activity, and might also be of potential use for future treatment strategies of sciatica.

After previously being considered as just a biologically inactive tissue component compressing the spinal nerve root at disc herniation, the nucleus pulposus has recently been found to be highly active, inducing both structural and functional changes in adjacent nerve roots when applied epidurally (24,37,38,41,42). It has thereby been established that autologous nucleus pulposus may induce axonal changes and a characteristic myelin injury (24,38,41,42), increased vascular permeability (9,44), intra vascular coagulation (24,36), and that membrane-bound structure or substances of the nucleus pulposus-cells are responsible for these effects (24,37). The effects have also been found to be efficiently blocked by methyl-prednisolone and cyclosporin A (2,38). When critically looking at these data, one realizes that there is at least one cytokine that relates to all of these effects, Tumor Necrosis Factor alpha (TNF- α). To assess if TNF- α may be involved in the nucleus pulposus-induced nerve root injury the presence of TNF- α in nucleus pulposus-cells was assessed and was studied if the nucleus pulposus-induced effects could be blocked by doxycycline and also by a selective monoclonal antibody.

MATERIAL AND METHODS

Series-1, Presence of TNF- α in pig nucleus pulposus-cells:

Nucleus pulposus (NP) from a total of 13 lumbar and thoracic discs were obtained from a pig used for other purposes. NP was washed once in Ham's F12 medium (Gibco BRL, Paisley, Scotland) and then centrifuged and suspended in 5 ml of collagenase solution in Ham's F12 medium (0.8 mg/ml, Sigma Chemical Co., St Louis, MO, USA) for 40 minutes, at 37°C in 25 cm² tissue culture flasks. The separated NP-cell pellets were suspended in DMEM/F12 1:1 medium (Gibco BRL, Paisley, Scotland) supplemented with 1% L-glutamine 200 mM (Gibco BRL, Paisley, Scotland), 50 µg/ml gentamycine sulphate (Gibco BRL, Paisley, Scotland) and 10% foetal calf serum (FCS), (Gibco BRL, Paisley, Scotland). The cells were cultured at 37°C and 5% CO₂ in air for 3-4 weeks and then cultured directly on tissue culture treated glass slides (Becton Dickinson & Co Labware, Franklin Lakes, NJ, USA). After 5 days on the glass slides, the cells were fixed in situ by acetone for 10 minutes. After blocking irrelevant antigens by application of 3% H₂O₂ (Sigma Chemical Co., St Louis, MO, USA) for 30 minutes and Horse Serum (ImmunoPure ABC, peroxidase

mouse IgG staining kit nr.32028, Pierce, Rockford, IL) for 20 minutes, the primary antibody (Anti-pig TNF- α monoclonal purified antibody, Endogen, Cambridge, MA, USA) was applied over night at +40°C, diluted at 1:10, 1:20 and 1:40. For control, BSA (bovine serum albumin, Intergen Co, New York, USA) suspended in PBS (phosphate buffered saline, Merck, Darmstadt, Germany) was applied in the same fashion. The next day the cells were washed with 1% BSA in PBS and the secondary antibody (ImmunoPure ABC, peroxidase mouse IgG staining kit nr.32028, Pierce, Rockford, IL) was applied for 30 minutes. To

enhance this reaction, the cells were exposed to Avidin-Biotin complex for additionally 30 minutes (ImmunoPure ABC, peroxidase mouse IgG staining kit nr.32028, Pierce, Rockford, IL). The cells were then exposed to 20 mg of DAB (3,3-diaminobenzidine tetrahydrochloride nr. D-5905, Sigma Chemical Co., St Louis, MO, USA) and 0.033 ml of 3% H₂O₂ in 10 ml of saline for 10 minutes. The cells were washed in PBS, dehydrated in a series of ethanol, mounted and examined by light microscopy by an unbiased observer regarding the presence of a brown colouration indicating presence of TNF- α .

Series-2, Neurophysiologic evaluation:

Thirteen pigs, (body weight 25-30 kg) received an intramuscular injection of 20 mg/kg body weight of Ketalar^R (ketamine 50 mg/ml, Parke-Davis, Morris Plains, New Jersey) and an intravenous injection of 4 mg/kg body weight of Hypnodil^R (methomidate chloride 50 mg/ml, AB Leo, Helsingborg, Sweden) and 0.1 mg/kg body weight of Stresnil^R (azaperon 2 mg/ml, Janssen Pharmaceutica, Beerse, Belgium). Anaesthesia was maintained by additional intravenous injections of 2 mg/kg body weight of Hypnodil^R and 0.05 mg/kg body weight of Stresnil^R. The pigs also received an intravenous injection of 0.1 mg/kg of Stesolid Novum^R (Diazepam, Dumex, Helsingborg, Sweden) after surgery.

Nucleus pulposus was harvested from the 5th lumbar disc through a retro peritoneal approach (42). Approximately 40 mg of the nucleus pulposus was applied to the

sacroccygeal cauda equina through a midline incision and laminectomy of the first coccygeal vertebra. Four pigs did not receive any treatment (no treatment). Four other pigs received an intravenous infusion of 100 mg of doxycycline (Vibramycino, Pfizer Inc., New York, USA) in 100 ml of saline over 1 hour. In 5 pigs, the nucleus pulposus was mixed with

100 gl of a 1,11 mg/ml suspension of the anti-TNF- α antibody used in series 1, before application.

5 Three days after the application, the pigs were reanaesthetized by an intramuscular injection of 20 mg/kg body weight of Ketalar^R and an intravenous injection of 35 mg/kg body weight of Pentothal^R (Thiopental sodium, Abbott lab, Chicago, IL). The pigs were ventilated on a respirator. Anaesthesia was maintained by an intravenous bolus injection of 100 mg/kg

10 body weight of Chloralose (α)-D(+)-gluco-chloralose, Merck, Darmstadt, Germany) and by a continuous supply of 30 mg/kg/hour of Chloralose. A laminectomy from the 4th sacral to the 3rd coccygeal vertebra was performed. The nerve roots were covered with Spongostane^R (Ferrosan, Demnark). Local tissue temperature was continuously monitored and maintained at 37.5-38.0°C by means of a heating lamp.

15 The cauda equina was stimulated by two E2 subdermal platinum needle electrodes (Grass Instrument Co., Quincy, MA) which were connected to a Grass SD9 stimulator (Grass Instrument Co., Quincy, MA) and gently placed intermittently on the cauda equina first 10 mm cranial and then 10 mm caudal to the exposed area. To ensure that only impulses from exposed nerve fibers were registered, the nerve root that exited from the spinal canal between the two stimulation sites were cut. An EMG was registered by two subdermal

20 platinum needle electrodes which were placed into the paraspinal muscles in the tail approximately 10 mm apart. This procedure is reproducible and represents a functional measurement of the motor nerve fibers of the cauda equina nerve roots. The EMG was visualized using a Macintosh IICI computer provided with Superscope software and MacAdios II AID converter (GW Instruments, Sommerville, MA) together with a Grass P18

25 preamplifier (Grass Instrument Co., Quincy, MA). The separation distance between the first peaks of the EMG from the two recordings was determined and the separation distance between the two stimulation sites on the cauda equina was measured with calipers. The

30 nerve conduction velocity between the two stimulation sites could thus be calculated from these two measurements.

The person performing the neurophysiologic analyses was unaware of the experimental protocol for the individual animal, and after finishing the complete study the data were

arranged in the three experimental groups and statistical differences between the groups were assessed by Student's t-test. The experimental protocol for this experiment was approved by the local animal research ethics committee.

RESULTS

5 Series-1, Presence of TNF- α in pig nucleus pulposus-cells.

Examples of the light microscopic appearance of the stained glass slides. In the sections using BSA in PBS as "primary antibody" (control) no staining was observed, ensuring that there was no labelling and visualization of irrelevant antigens. When the anti-TNF- α antibody was applied at 1:40 dilution there was only a weak staining. However, the staining increased with diminishing dilutions of the antibody. The staining was seen in the soma of the cells and it was not possible to differentiate whether TNF- α was located in the cytoplasm, on the cell surface bound to the cell-membrane, or both.

10 Series-2, Neurophysiologic evaluation:

Application of non-modified nucleus pulposus and without any treatment induced a reduction in nerve conduction velocity similar to previous studies (no treatment, figure 2), whereas treatment with doxycycline completely blocked this reduction ($p < 0.01$ Student's t-test). Local application of anti-TNF- α -antibody also induced a partial block of this reduction, although not as complete as doxycycline and not statistically significant to the no treatment-series.

20 DISCUSSION

The data of the present study demonstrated that TNF- α may be found in nucleus pulposus-cells of the pig. If TNF- α was blocked by a selective monoclonal antibody, the nucleus pulposus-induced reduction of nerve root conduction velocity was partially blocked, although no statistically significant compared to the series with non-treated animals.

25 However, if doxycycline was used to inhibit TNF- α , the reduction of nerve conduction velocity was significantly blocked.

30 In recent years, it has been verified that local application of autologous nucleus pulposus may injure the adjacent nerve roots. Thus, it has become evident that the nerve root injury seen at disc herniation may not be solely based on mechanical deformation of the nerve root, but may also be induced by unknown "biochemical effects" related to the epidural presence

of herniated nucleus pulposus. Although this new research field has generated many experimental studies, the mechanisms and substances involved are not fully known. It has been seen that local application of autologous nucleus pulposus may induce axonal injury (24,37,38,40-42), a characteristic injury of the myelin sheath (24,38,40-42), a local increase of vascular permeability (9,36,44), intra vascular coagulations, reduction of intra neural blood flow (43), and leukotaxis (36). It has been seen that the nucleus pulposus-related effects may be blocked efficiently by methylprednisolone (38) and cyclosporin A (2), and slightly less efficiently by indomethacin (3), and lidocaine (69). Further, it has been understood that the effects are mediated by the nucleus pulposus-cells (37), particularly by substances or structures bound to the cell-membranes (25). When critically considering these data, it becomes evident that at least one specific cytokine could be related to these observed effects, Tumor Necrosis Factor-alpha (TNF- α). TNF- α may induce nerve injury (29,31,45,50,66) mainly seen as a characteristic myelin injury that closely resembles the nucleus pulposus-induced myelin-injury (29,47,51,54,62,64,66,70). TNF- α may also induce an increase in vascular permeability (47,66) and initiate coagulation (22,34,63). Further, TNF- α may be blocked by steroids (4,8,21,61,68), and cyclosporin A (11,55,67,68). However, the blocking effect on TNF- α is not so pronounced by NSAID (14,17,20) and very low or the opposite by lidocaine (5,32,46,60). It was recently observed that local application of nucleus pulposus may induce pain-related behaviour in rats, particularly thermal hyperalgesia (23,40). TNF- α has also been found to be related to such pain-behaviouristic changes (12,35,56,66), and also to neuropathies in general (30,54,56,57). However there are no studies that have assessed the possible presence of TNF- α in the cells of the nucleus pulposus.

To assess if TNF- α could be related to the observed nucleus pulposus induced reduction in nerve root conduction velocity it was necessary first to analyse if there were TNF- α in the nucleus pulposus-cells. The data clearly demonstrated that TNF- α was present in these cells.

TNF- α is produced as precursor (pro-TNF) that is bound to the membrane and it is activated by cleavage from the cell-membrane by a zinc-dependent metallo-endopeptidase (TNF-alpha converting enzyme, TACE) (6,15,16,48,49). This may thus relate well to experimental findings where application of the mere cell-membranes of autologous nucleus pulposus-cells

induced nerve conduction velocity reduction, which indicated that the effects were mediated by a membrane-bound substances. Second, the effects of the TNF- α had to be blocked in a controlled manner. We then first choose to add the same selective antibody that was used for immunohistochemistry in series 1, which is known to also block the effects of TNF- α , to the nucleus pulposus before application. Also, we choose to treat the pigs with doxycycline, which is known to block TNF- α (26,27,33,52,53). However, due to the low pH of the doxycycline preparation it was chosen to treat the pigs by intravenous injection instead of local addition to the nucleus pulposus since nucleus pulposus at a low pH has been found to potentiate the effects of the nucleus pulposus (38,39).

The data regarding nerve conduction velocity showed that the reduction was completely blocked by the doxycycline-treatment and that the nerve conduction velocity in this series was close to the conduction velocity after application of a control substance (retro peritoneal fat) from a previous study (42). Application of the anti-TNF- α -antibody to the nucleus pulposus also partially blocked the reduction in velocity, however, not as pronounced as doxycycline, and the velocity in this series was not statistically different to the velocity in the series with not treated animals, due to the wide deviation of the data.

The fact that the anti-TNF- α antibody treatment only partially blocked the nucleus pulposus-induced reduction of nerve conduction velocity and the high standard deviation of the data could probably have at least two different explanations. First, if looking at the specific data within this group it was found that the nerve conduction velocity was low in 2 animals (mean 37.5 m/s) and high in 3 animals (mean 81.3 m/s). There are thus 2 groups of distinctly different data within the anti-TNF- α treatment series. This will account for the high standard deviation and might imply that the blocking effect was sufficient in 3 animals and non-sufficient in 2 animals. The lack of effects in these animals could be based simply on the amount of antibodies in relation to TNF- α molecules not being sufficient, and if a higher dose of the antibody had been used, the TNF- α effects would thus have been blocked even in these animals. Such a scenario could then theoretically imply that TNF- α alone is responsible for the observed nucleus pulposus-induced effects, and that this could not be verified experimentally due to the amount of antibody being too low.

Second, it is also known that tetracyclines such as doxycycline and minocycline may block a number of cytokines and other substances. For instance they may block IL-1 (1,28,58), IFN γ (27), NO-synthetasel, and metalloproteinases (1,53,58). Particularly IL-1 and IFN γ are known to act synergistically with TNF- α and are known to be more or less neurotoxic (7,10,13,18,19,56,59). These substances are also blocked by steroids and cyclosporin A which corresponds well with the previous observations on nucleus pulposus-induced nerve root injury which have shown that the nucleus pulposus-induced effects may be blocked by these substances (8,67). One may therefore also consider the possibility that a selective block of TNF- α may not be sufficient to completely block the nucleus pulposus-induced effects on nerve function, and that simultaneous block of other synergistic substances is necessary as well. Thus, this scenario, on the other hand, implies that TNF- α is not solely responsible for the nucleus pulposus-induced effects, and that other synergistic substances, which are also blocked by doxycycline, may be necessary.

TNF- α may have various pathophysiologic effects. It may have direct effects on tissues such as nerve tissue and blood vessels, it may trigger other cells to produce other pathogenic substances and it may trigger release of more TNF- α both by inflammatory cells and also by Schwann-cells locally in the nerve tissue (65). There is thus reason to believe that even low amounts of TNF- α may be sufficient to initiate these processes and that there is a local recruitment of cytokine producing cells and a subsequent increase in production and release of other cytokines as well as TNF- α . TNF- α may therefore act as the "ignition key" of the pathophysiologic processes and play an important role for the initiation of the pathophysiologic cascade behind the nucleus pulposus-induced nerve injury.

In conclusion, although the exact role of TNF- α can not be fully understood from the experimental set-up, we may conclude that for the first time a specific substance (TNF- α) has been linked to the nucleus pulposus-induced nerve root injury, probably potentiated by other substances which are also blocked by doxycycline, such as for instance IL1, IFN γ and NO-synthetase. This new information may be of significant importance for the continued understanding of nucleus pulposus-induced nerve injury as well as raising the question of the potential future clinical use of pharmacological interference with TNF- α and related

1998 -09- 2 5

Huvudfaxen Kassan

11

substances, by for instance doxycycline or metalloproteinase-inhibitors, for the treatment of sciatica.

5 The presence of TNF- α in pig nucleus pulposus-cells was thus immunohistochemically verified. Block of TNF- α by a monoclonal antibody partially limited the nucleus pulposus-induced reduction of nerve root conduction velocity, whereas intravenous treatment with doxycycline significantly blocked this reduction. These data for the first time links one specific substance, TNF- α , to the nucleus pulposus-induced nerve injury.

10 The compounds of the invention can be administered in a variety of dosage forms, e.g., orally, in the form of tablets, capsules, sugar or film coated tablets, liquid solutions; rectally, in the form of suppositories; parenterally, e.g., intramuscularly or by intravenous injection or infusion. The therapeutic regimen for the different clinical syndromes must be adapted to the type of pathology taken in to account, as usual, also the route of administration, the form in
15 which the compound is administered and age, weight, and condition of the subject involved.

20 The oral route is employed, in general, for all conditions, requiring such compounds. In emergency cases preference is given to intravenous injection. For these purposes the compounds of the invention can be administered orally at doses ranging from about 20 to about 1500 mg/day. Of course, these dosage regimens may be adjusted to provide the optimal therapeutic response.

25 The nature of the pharmaceutical composition containing the compounds of the invention in association with pharmaceutically acceptable carriers or diluents will, of course, depend upon the desired route of administration. The composition may be formulated in the conventional manner with the usual ingredients. For example, the compounds of the invention may be administered in the form of aqueous or oily solutions or suspensions, tablets, pills, gelatine capsules (hard or soft ones) syrups, drops or suppositories.

30 Thus for oral administration, the pharmaceutical compositions containing the compounds of the invention are preferably tablets, pills or gelatine capsules, which contain the active substance together with diluents, such as lactose, dextrose, sucrose, mannitol, sorbitol,

cellulose; lubricants, e.g., silica, talc, stearic acid, magnesium or calcium stearate, and/or polyethylene glycols; or they may also contain binders, such as starches, gelatine, methyl cellulose, carboxymethylcellulose, gum arabic, tragacanth, polyvinylpyrrolidone; disaggregating agents such as starches, alginic acid, alginates, sodium starch glycolate, microcrystalline cellulose; effervescing agents such as carbonates and acids; dyestuffs; sweeteners; wetting agents, such as lecithin, polysorbates, laurylsulphates; and in general non-toxic and pharmaceutically inert substances used in the formulation of pharmaceutical

compositions. Said pharmaceutical compositions may be manufactured in known manners, e.g., by means of mixing, granulating, tableting, sugar-coating or film-coating processes. In the case film providing compounds can be selected to provide release in the right place in the intestinal tract with regard to absorption and maximum effect. Thus pH-dependent film formers can be used to allow absorption in the intestines as such, whereby different phthalates are normally used or acrylic acid/methacrylic acid derivatives and polymers.

The liquid dispersions for oral administration may be e.g., syrups, emulsion, and suspensions.

The syrups may contain as carrier, e.g., saccharose, or saccharose with glycerine and/or mannitol and/or sorbitol.

Suspensions and emulsions may contain as carrier, e.g., a natural gum, such as gum arabic, xanthan gum, agar, sodium alginate, pectin, methyl cellulose, carboxymethylcellulose, polyvinyl alcohol.

The suspension or solutions for intramuscular injections may contain together with the active compound, a pharmaceutically acceptable carrier, such as e.g., sterile water, olive oil, ethyl oleate, glycols, e.g., propylene glycol, and if so desired, a suitable amount of lidocaine hydrochloride. Adjuvants for triggering the injection effect can be added as well.

The solutions for intravenous injection or infusion may contain as carrier, e.g., sterile water, or preferably, a sterile isotonic saline solution.

1998 -09- 25

Huvudtaxan Kassan

13

The suppositories may contain together with the active compound, a pharmaceutically acceptable carrier, e.g., cocoa-butter polyethylene glycol, a polyethylene sorbitan fatty acid ester surfactant or lecithin.

REFERENCES

1. Amin AR, Attur MG, Thakker GD, Patel PD, Vyas PR, Patel RN, Patel IR, Abramson SB. A novel mechanism of action of tetracyclines: effects on nitric oxide syntheses. Proc Natl Acad Sci U S A 1996; 93:14014-9.
2. Arai I, Konno S, Otani K, Kikuchi S, Olmarker K. Cyclosporin A blocks the toxic effects of nucleus pulposus on spinal nerve roots. Manuscript
3. Arai I, Mao GP, Otani K, Konno S, Kikuchi S, Olmarker K. Indomethacin blocks nucleus pulposus related effects in adjacent nerve roots. Manuscript
4. Baumgartner RA, Deramo VA, Beaven MA. Constitutive and inducible mechanisms for synthesis and release of cytokines in immune cell lines. J Immunol 1996;157:4087-93.
5. Bidani A, Heming TA. Effects of lidocaine on cytosolic pH regulation and stimulus-induced effector functions in alveolar macrophages. Lung 1997;175:349-61.
6. Black RA, Rauch CT, Kozlosky CJ, Peschon JJ, Slack JL, Wolfson MF, Castner BJ, Stocking KL, Reddy P, Srinivasan S, Nelson N, Boiani N, Schooley KA, Gerhart M, Davis R, Fitzner JN, Johnson RS, Paxton RJ, March CJ, Cerretti DP. A metalloproteinase disintegrin that releases tumour-necrosis factor- alpha from cells. Nature 1997;385:729-33.
7. Bluthé RM, Dantzer R, Kelley KW. Interleukin-1 mediates behavioural but not metabolic effects of tumor necrosis factor alpha in mice. Eur J Pharmacol 1991;209:281-3.
8. Brattsand R, Linden M. Cytokine modulation by glucocorticoids: mechanisms and actions in cellular studies. Aliment Pharmacol Ther 1996; 10:81-90.
9. Byrdd G, Otani K, Rydevik B, Olmarker K. Acute increase in endoneural vascular permeability induce by epidural application of nucleus pulposus on spinal nerve roots. Manuscript
10. Chao CC, Hu S, Ehrlich L, Peterson PK. Interleukin-1 and tumor necrosis factor-alpha synergistically mediate neurotoxicity: involvement of nitric oxide and of N-methyl-D-aspartate receptors. Brain Behav Immun 1995;9:355-65.
11. Dawson J, Hurtenbach U, MacKenzie A. Cyclosporin A inhibits the in vivo production of interleukin-1beta and tumour necrosis factor alpha, but not interleukin-6, by a T-cell-independent mechanism. Cytokine 1996;8:882-8.

12. DeLeo JA, Colburn RW, Rickman AJ. Cytokine and growth factor immunohistochemical spinal profiles in two animal models of mononeuropathy. Brain Res 1997;759:50-7.

13. Gadiant RA, Cron KC, Otten U. Interleukin-1 beta and tumor necrosis factor-alpha synergistically stimulate nerve growth factor (NGF) release from cultured rat astrocytes. Neurosci Lett 1990;117:335-40.

14. Garcia-Vicuna R, Diaz-Gonzalez F, Gonzalez-Alvaro I, del Pozo MA, Moilinedo F, Cabanas C, Gonzalez-Amaro R, Sanchez-Madrid F. Prevention of cytokine-induced changes in leucocyte adhesion receptors by nonsteroidal antiinflammatory drugs from the oxicam family. Arthritis Rheum 1997;40:143-53.

15. Gearing AJ, Beckett P, Christodoulou M, Churchill M, Clements J, Davidson AH, Drummond AH, Galloway WA, Gilbert R, Gordon JL, et al. Processing of tumour necrosis factor-alpha precursor by metalloproteinases. Nature 1994;370:555-7.

16. Gazelle EJ, Banda MJ, Leppert D. Matrix metallo-proteinases in immunity. J Immunol 1996; 156: 14.

17. Gonzalez E, de la Cruz C, de Nicolas R, Egido J, Herrero-Beaumont G. Long-term effect of nonsteroidal anti-inflammatory drugs on the production of cytokines and other inflammatory mediators by blood cells of patients with osteosis. Agents Actions 1994;41:171-8.

18. Hartung HP, Jung S, Stoll G, Zielasek J, Schmidt B, Archelos JJ, Toyka KV. Inflammatory mediators in demyelinating disorders of the CNS and PNS. J Neuroimmunol 1992;40:197-210.

19. Hattori A, Iwasaki S, Murase K, Tsujimoto M, Sato M, Hayashi K, Kohno M. Tumor necrosis factor is markedly synergistic with interleukin 1 and interferon-gamma in stimulating the production of nerve growth factor in fibroblasts. FEBS Lett 1994;340:177-80.

20. Herman JH, Sowder WG, Hess EV. Nonsteroidal antiinflammatory drug modulation of prosthesis pseudomembrane induced bone resorption. J Rheumatol 1994;21:338-43.

21. Iwamoto S, Takeda K. [Possible cytotoxic mechanisms of TNF in vitro]. Hum Cell 1990;3:107-12.

22. Jurd KM, Stephens CJ, Black MM, Hunt BJ. Endothelial cell activation in cutaneous

vasculitis. Clin Exp Dermatol 1996;21:28-32.

23. Kawakami M, Tamaki T, Weinstein JN, Hashizume H, Nishi H, Meller ST.

Pathomechanism of pain-related behaviour produced by allografts of intervertebral disc in the rat. Spine 1996;21:2101-7.

- 5 24. Kayama S, Konno S, Olmarker K, Yabuki S, Kikuchi S. Incision of the anulus fibrosis induces nerve root morphologic, vascular, and functional changes. An experimental study. Spine 1996;21:2539-43.

10 25. Kayama S, Olmarker K, Larsson K, Sjögren-Jansson E, Lindahl A, Rydevik B. Cultured, autologous nucleus pulposus cells induce structural and functional changes in spinal nerve roots. Manuscript

26. Kloppenburg M, B-an BM, de Rooij-Dijk HH, Miltenburg AM, Daha MR, Breedveld FC,

15 Dijkmans BA, Verweij C. The tetracycline derivative minocycline differentially affects cytokine production by monocytes and T lymphocytes. Antimicrob Agents Chemother 1996;40:934-40.

27. Kloppenburg M, Verweij CL, Miltenburg AM, Verhoeven AJ, Daha MR, Dijkmans BA, Breeveld FC. The influence of tetracyclines on T cell activation. Clin Exp Immunol 1995;102:635-41.

20 28. Lamster IB, Pullman JR, Celenti RS, Grbic JT. The effect of tetracycline fiber therapy on beta-glucuronidase and interleukin-1 beta in crevicular fluid. J Clin Periodontol 1996;23:816-22.

29. Liberski PP, Yanagihara R, Nerurkar V, Gajdusek DC. Further ultrastructural studies of lesions induced in the optic nerve by tumor necrosis factor alpha (TNF-alpha): a comparison with experimental Creutzfeldt-Jakob disease. Acta Neurobiol Exp (Warsz) 1994;54:209-18.

25 30. Lin XH, Kashima Y, Khan M, Heller KB, Gu XZ, Sadun AA. An immunohistochemical study of TNF-alpha in optic nerves from AIDS patients. Curr Eye Res 1997;16:1064-8.

31. Madigan MC, Sadun AA, Rao NS, Dugel PU, Tenhula WN, Gill PS. Tumor necrosis factor-alpha (TNF-alpha)-induced optic neuropathy in rabbits. Neurol Res 1996; 18:176-84.

30 32. Matsumori A, Ono K, Nishio R, Nose Y, Sasayama S. Amiodarone inhibits production of tumor necrosis factor-alpha by human mononuclear cells: a possible mechanism for its effect in heart failure. Circulation 1997;96:1386-9.

33. Milano S, Arcoleo F, D'Agostino P, Cillari E. Intraperitoneal injection of tetracyclines protects mice from lethal endotoxemia downregulating inducible nitric oxide synthase in various organs and cytokine and nitrate secretion in blood. Antimicrob Agents Chemother 1997;41:117-21.

5 34. Nawroth P, Handley D, Matsueda G, De Waal R, Gerlach H, Blohm D, Stem D. Tumor necrosis factor/cachectin-induced intra vascular fibrin formation in meth A fibrosarcomas. J Exp Med 1988;168:637-47.

10 35. Oka T, Wakugawa Y, Hosoi M, Oka K, Hori T. Intracerebroventricular injection of tumor necrosis factor-alpha induces thermal hyperalgesia in rats. Neuroimmunomodulation 1996;3:135-40.

36. Olmarker K, Blomquist J, Stromberg J, Nannmark, U, Thomsen P, Rydevik B. Inflammation-togenic properties of nucleus pulposus. Spine 1995;20:665-9.

15 37. Olmarker K, Brisby H, Yabuki S, Nordborg C, Rydevik B. The effects of normal, frozen, and hyaluronidase-digested nucleus pulposus on nerve root structure and function. Spine 1997;22:4715; discussion 476.

38. Olmarker K, Byrod G, Comesfjord M, Nordborg C, Rydevik B. Effects of methylprednisolone on nucleus pulposus-induced nerve root injury. Spine 1994; 19:1803-8.

39. Olmarker K, Iwabuchi M, Larsson K, Rydevik B. Effects of in vitro degenerated nucleus pulposus on nerve root conduction velocity. Manuscript

20 40. Olmarker K, Myers RR. Pathogenesis of sciatic pain: Role of herniated nucleus pulposus and deformation of spinal nerve root and DRG. Manuscript

41. Olmarker K, Nordborg C, Larsson K, Rydevik B. Ultrastructural changes in spinal nerve roots induced by autologous nucleus pulposus. Spine 1996;21:411-4.

25 42. Olmarker K, Rydevik B, Nordhorn C. Autologous nucleus pulposus induces neurophysiologic and histologic changes in porcine cauda equina nerve roots [see comments]. Spine 1993;18:1425-32.

43. Otani K, Arai I, Mao GP, Konno S, Olmarker K, Kikuchi S. Nucleus pulposus-induced nerve root injury. The relationship between blood flow and nerve conduction velocity. Manuscript

30 44. Otani K, Mao GP, Arai I, Konno S, Olmarker K, Kikuchi S. Nucleus pulposus-induced increase in vascular permeability in the nerve root. Manuscript

45. Petrovich MS, Hsu HY, Gu X, Dugal P, Heller KB, Sadun AA. Pentoxifylline suppression of TNF- alpha mediated axonal degeneration in the rabbit optic nerve. Neurol Res 1997; 19:551-4.

46. Pichler WJ, Zanni M, von Greyerz S, Schnyder B, Mauri-HeUweg D, Wendland, T.
5 High IL-5
production by human drug-specific T cell clones. Int Arch Allergy Immunol 1997; 1 13 :177-80.

47. Redford EJ, Hall SM, Smith KJ. Vascular changes and demyelination induced by the intra neural injection of tumour necrosis factor. Brain 1995; 1 18 :869-78.

10 48. Robache-Gallea S, Bruneau JM, Robbe H, Morand V, Capdevila C, Bhatnagar N, Chouaib S, Roman-Roman S. Partial purification and characterization of a tumor necrosis factor- alpha converting activity. Eur J Immunol 1997;27:1275-82.

49. Rosendahl MS, Ko SC, Long DL, Brewer MT, Rosenzweig B, Hedl E, Anderson L, Pyle SM, Moreland J, Meyers MA, Kohno T, Lyons D, Lichenstein HS. Identification and
15 characterization of a pro-tumor necrosis factor- alpha-processing enzyme from the ADAM family of zinc metalloproteases. J Biol Chem 1997;272:24588-93.

50. Said G, Hontebeyrie-Joskowicz M. Nerve lesions induced by macrophage activation. Res Immunol 1992;143:589-99.

20 51. Sehnaj KW, Raine CS. Tumor necrosis factor mediates myelin and oligodendrocyte damage in vitro. Ann Neurol 1988;23:339-46.

52. Shapira L, Houri Y, Barak V, Halabi A, Soskoine WA, Stabholz A. Human monocyte response to cementum extracts from periodontally diseased teeth: effect of conditioning with tetracycline. J Periodontol 1996;67:682-7.

53. Shapira L, Houri Y, Barak V, Soskolne WA, Halabi A, Stabholz A. Tetracycline inhibits
25 Porphyromonas gingivalis lipopolysaccharide- induced lesions in vivo and TNF alpha processing in vitro. J Periodontal Res 1997;32:183-8.

54. Sharief MK, Ingram DA, Swash M. Circulating tumor necrosis factor-alpha correlates with electrodiagnostic abnormalities in Guillain-Barre syndrome. Ann Neurol 1997;42:68-73.

30 55. Smith CS, Ortega G, Parker L, Shearer WT. Cyclosporin A blocks induction of tumor necrosis factor-alpha in human B lymphocytes. Biochem Biophys Res Commun

1994;204:383-90.

56. Sonuner C, Schmidt C, George A, Toyka KV. A metalloprotease-inhibitor reduces pain associated behaviour in mice with experimental neuropathy. Neurosci Lett 1997;237:45-8.

57. Sorkin LS, Xiao WH, Wagner R, Myers RR. Tumour necrosis factor-alpha induces ectopic activity in nociceptive primary afferent fibres. Neuroscience 1997;81:255-62.

58. Steinmeyer J, Daufeldt S, Taiwo YO. Pharmacological effect of tetracyclines on proteoglycanases from interleukin-1-treated articular cartilage. Biochem Pharmacol 1998;55:93-100.

59. Stoll G, Jung S, Jander S, van der Meide P, Hartung HP. Tumor necrosis factor-alpha in immunomediated demyelination and Wallerian degeneration of the rat peripheral nervous system. Neuroimmunol 1993;45:175-82.

60. Takao Y, Mikawa K, Nishina K, Maekawa N, Obara H. Lidocaine attenuates hyperoxic lung injury in rabbits. Acta Anaesthesiol Scand 1996;40:318-25.

61. Teoh KH, Bradley CA, Gauldie J, Burrows H. Steroid inhibition of cytokine-mediated vasodilation after warm heart surgery. Circulation 1995;92:II347-53.

62. Tsukamoto T, Ishikawa M, Yamamoto T. Suppressive effects of TNF-alpha on myelin formation in vitro. Acta Neurol Scand 1995;91:71-5.

63. van der Poll T, Jansen PM, Van Zee KJ, Welborn MBr, de Jong I, Hack CE, Loetscher H, Lesslauer W, Lowry SF, Moidawer LL. Tumor necrosis factor-alpha induces activation of coagulation and fibrinolysis in baboons through an exclusive effect on the p55 receptor. Blood 1996;88:922-7.

64. Villarroja H, Violleau K, Ben Younes-Chennoufi A, Baumann N. Myelin-induced experimental allergic encephalomyelitis in Lewis rats: tumor necrosis factor alpha levels in serum of cerebrospinal fluid immunohistochemical expression in glial cells and neurophages of optic nerve and spinal cord. J Neuroimmunol 1996;64:55-61.

65. Wagner R, Myers RR. Schwann cells produce tumor necrosis factor alpha: expression in injured non-injured nerves. Neuroscience 1996;73:625-9.

66. Wagner R, Myers RR. Endoneurial injection of TNF-alpha produces neuropathic pain behaviours. Neuroreport 1996;7:2897-901.

67. Wasaki S, Sakaida I, Uchida K, Kiinura T, Kayano K, Okita K. Preventive effect of cyclosporin A on experimentally induced acute liver injury in rats. Liver 1997; 17:107-14.

Ink. t. Patent- och reg.verket

1998 -09- 25

Huvudfaxen Kassa

20

68. Wershil BK, Furuta GT, Lavigne JA, Choudhury AR, Wang ZS, Galli SJ.

Dexamethasone cyclosporin A suppress mast cell-leukocyte cytokine cascades by multiple mechanisms. Int Arch Allergy Immunol 1995;107:323-4.

69. Yabuki S, Kawaguchi Y, Olmarker K, Rydevik B. Effects of lidocaine on nucleus
5 pulposus-induced nerve root injury. Manuscript

70. Zhu J, Bai XF, Mix E, Link H. Cytokine dichotomy in peripheral nervous system
influences the outcome of experimental allergic neuritis: dynamics of mRNA expression for
IL-1 beta, IL-6, IL-10, IL-12, TNF-alpha, TNF-beta, and cytolytic. Clin Immunol
Immunopathol 1997;84:85-94.

10

CLAIMS

1. Pharmaceutical composition for the inhibition of spinal disk TNF- α for the treatment of nerve root injury comprising a pharmaceutically effective amount of a metallo proteinase inhibitor together with a pharmaceutically acceptable carrier.

5

2. Pharmaceutical composition according to claim 1, wherein the active component is selected from the group consisting of tetracycline, doxycycline, lymecycline, oxytetracycline, minocycline, dedimethylaminotetracycline, hydroxamic acid compounds, carbocyclic acids and derivatives, cyclosporine, methylprenisolone, thalidomide, lazaroides, pentoxiphylline in the form of bases or addition salts.

10

3. Pharmaceutical composition according to claim 2, wherein the active component is doxycycline.

15

4. Method for the treatment of spinal disorders as nerve root injury caused by the liberation of TNF- α in mammals, including man, comprising the administration of a pharmaceutically effective amount of a metallo proteinase inhibitor.

20

5. Use of a metallo proteinase inhibitor in the preparation of a pharmaceutical composition for the treatment of spinal disorders as nerve root injury caused by the liberation of TNF- α .

Ink. t. Patent- och reg.verkst

1998 -09- 25

Huvudfaxen Kassa

22

ABSTRACT

The present invention relates to pharmaceutical compositions for the treatment of spinal disorders caused by the liberation of TNF- α comprising an effective amount of a metallo-proteinase inhibitor, as well as method for treatment of such disorders, and the use of metallo- proteinase inhibitors in the preparation of pharmaceutical composition for such treatment.

5

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.